

Predictors of HIV prevention knowledge and sexual behaviors among students at Makerere University Kampala, Uganda

Afra Nuwasiima^(1,2), Nelson Owuor Onyango⁽¹⁾, Patricia Navvuga⁽²⁾, Elly Nuwamaya⁽²⁾, Agnes Nyabigambo⁽³⁾, Solomon J. Lubinga⁽⁴⁾, Joseph B. Babigumira⁽⁴⁾

(1) School of Mathematics, University of Nairobi, Kenya

(2) Global Health Economics, Kampala, Uganda

(3) Department of Community Health and Behavioral Sciences, School of Public Health, Makerere University Kampala, Uganda

(4) Department of Global Health, University of Washington, Seattle, WA, USA

CORRESPONDING AUTHOR: Afra Nuwasiima, School of Mathematics, University of Nairobi, Kenya - Global Health Economics, Kampala, Uganda
email: anuwasiiima@gmail.com

DOI: 10.2427/12547

Accepted on August 29, 2017

ABSTRACT

Background: Prior reviews argue that unsafe sexual behaviors and poor HIV knowledge significantly increase the probability of acquiring HIV infections among adolescents. This study assessed the predictors of HIV prevention knowledge and sexual behaviors among Makerere University students in Uganda.

Methods: We performed a cross-sectional survey. We performed a normality test using Shapiro Wilk test on knowledge score. Results revealed that knowledge score was not normally distributed. The study used two sample Wilcoxon Rank Sum and Kruskal Wallis Rank tests to assess the effect of HIV knowledge on demographic characteristics and sexual behaviors. Post-hoc tests were conducted using Bonferroni correction. Spearman rank correlation test was used for continuous variables while Chi-square and Fisher's tests were used for categorical variables to assess the relationship between demographic characteristics and sexual behaviors. We used an iteratively re-weighted least square generalized linear model with an identity function to conduct multivariate regression analysis with knowledge score as the outcome variable.

Results: We report results for 1337 students. The mean age was 21.2SD (1.6) and more than half 700(52.4%) were male students. The median HIV prevention knowledge score of students was 13 IQR (11-15) in the range of 0 to 18. Males significantly scored higher than females (13.0 IQR (12-15) vs. 12.0 IQR (10-14) $p=0.000$), an increase in age was associated with higher knowledge scores ($Rho = 0.101$, $p = 0.000$). Students in the third year of study significantly scored higher than those in the first year, and government-sponsored students scored higher than the privately sponsored students. HIV knowledge was also significantly associated with sexual experience, and condom use at univariate level but insignificant at multiple level analysis. Males were more likely to have ever had sex (31.7% vs. 12.7%) and ever used a condom (63% vs. 55%) than females respectively

Conclusion: Our findings suggest that Makerere University students possessed good knowledge on HIV. There is evidence of an association between student's knowledge, and demographic characteristics and a few sexual behaviors. Future behavioral and educational programs that target both sexually and non-sexually experienced students should address the gender differences.

Key words: Sexual Health, HIV/AIDS, Knowledge, sexual behaviors, university students

BACKGROUND

Uganda started experiencing high rates of HIV infection in the late 1980s, and the infection reached its apex in the early 1990s [1]. The spread of HIV in the country is unevenly distributed among the population with women and urban dwellers at higher risk of infection [1]. The prevalence of HIV among Ugandans aged 15-49 increased from 6.4% in 2004 to 7.3 in 2011 [1,2]. Among the youth aged 20-29 years, the prevalence was estimated at 6.4% in 2011 [1]. It was estimated that in 2015, 4% of the country's population was living with HIV and 28,000 persons died of HIV-related illnesses [3].

Uganda is implementing various HIV/AIDS prevention programs that include; the comprehensive AIDS approach (Abstinence, Being faithful and Condom use), Prevention of Mother to Child Transmission (PMTCT), and voluntary medical male circumcision [1]. On the other hand, UNAIDS has listed five pillars for achieving a target of less than 500,000 new infections by 2020 like comprehensive sexuality education, economic empowerment and access to sexual and reproductive health services for young adults [3-5].

Earlier studies showed that 99% of Ugandans aged 15-49 years have ever heard about HIV and over 90% are aware of the mode of HIV transmission [1]. The proportion of comprehensive knowledge increased from 28 % in 2004 to 36% in 2011 among women aged 15-49 and from 36 % in 2004-05 to 43% in 2011 among men of the same age [1].

Adolescence is a crucial development step that may lead to the potential of unsafe sexual behaviors [6,7]. Risky lifestyles such as unprotected sex and inconsistent condom use are significantly associated with higher risk of HIV infections among adolescents [8-10]. Poor HIV/AIDS knowledge among the youth has also been indicated as a causal factor for increased HIV infections [5,11]. Studies demonstrated that university students are sexually adventurous, are often in multiple sexual relationships, and are less inclined to use condoms consistently [10,12].

Most HIV/AIDS prevention interventions in Uganda target the general population, and a few target high-risk populations, such as university students. Studies in Uganda have shown mixed evidence of the association between knowledge of HIV and unsafe sexual practices [13-17] [18-22]. We set out to assess the predictors of HIV knowledge on sexual behaviors among university students. The study adds to a growing body of literature on HIV prevention knowledge and sexual behaviors in similar settings and is one of the first endeavors to provide a quantitative score of knowledge on HIV in the country.

METHODS

Study design and setting

We performed a cross-sectional survey. The survey took place at Makerere University Kampala (MUK) among

students aged 18 to 30 years from March to April 2015. MUK is the largest university in Uganda and has a student population of over 4000. The University offers health services including HIV counseling and testing (HCT) to students in Uganda.

Sample size and sampling procedure

The sample size was calculated using the formula for a two-sample T-test. The study assumed a mean knowledge score difference of 0.7, a standard deviation $\delta_1 = \delta_2 = 4$, a 5% level of significance, and 90% power. A sample size of 686 was computed for each group leading to a total sample size of 1,372 participants. A sampling frame was generated from lists of university students obtained from the university where a random sample was taken. Eligibility of study participants included being a continuing student, aged between 18 to 30 years, possessing an Android smart-phone, and willing to provide written informed consent. Trained research assistants approached potential participants at hostels and screened them for inclusion, and then invited them to complete a self-administered tool in a private place. A total of 1,512 students completed the questionnaires. We report results of 1,337 students that provided complete information on the demographic characteristics and the outcome variable (HIV knowledge score).

Questionnaire

The questionnaire was developed and pre-tested before the main survey and contained three sections: socio-demographic characteristics; HIV prevention knowledge; and sexual and HIV prevention behavior.

The socio-demographic characteristics included questions on gender, age, university residence, home residence, marital status, year of study, religion, tuition status, and employment.

HIV/AIDS prevention knowledge was assessed using the 18 items version of the HIV knowledge questionnaire (KQ18) [23] to measure the student's knowledge. The HIV KQ18 is a brief self-administered measure of an individual's HIV-related knowledge and was designed to provide a more concise and well-established measure of HIV-related knowledge in street outreach, fieldwork, and intervention settings. The instrument contains 18 forced-choice statements ("true", "false", "don't know") related to knowledge on sexual transmission and the instrument yields a summary score by summing an individual's correct score across the 18 questions, with higher scores significant of greater HIV-related knowledge and with a scale of 0 to 18 [23].

The sexual and HIV prevention behaviors section of the questionnaire used questions modified from the Uganda AIDS Indicator Survey (UAIS). The UAIS is a nationally

representative, population-based, HIV serological survey implemented by Uganda Ministry of Health. The UAIS is designed to obtain national and subnational estimates of the prevalence of HIV and syphilis infection, data on attitudes towards HIV/AIDS, and sexual behavior. We collected data on: sexual intercourse, condom usage, alcohol use, sex with multiple partners, HIV testing, circumcision, willingness to circumcise, and knowledge of places where to acquire condoms.

Data collection and quality control

Data was collected using a pre-tested survey tool that was self-administered by the respondents. The trained interviewers helped to explain unfamiliar terminologies to the participants and made sure that students completed the required sections. Investigators went through the questionnaires daily to check for consistency and incompleteness.

Data management and statistical analysis

Data was double-entered and cross-validated to check for any errors in Epidata version 3.1. Data cleaning and analysis were conducted in Stata version 13.0 (Stata Corporation, College Station, Texas, USA) and R studio (open source version 1.0.136, R Foundation for Statistical Computing, Vienna, Austria). Descriptive analyses were performed using medians, and proportions. The study performed Shapiro Wilk test to check for normality. Normality tests indicated that the outcome variable (knowledge score) was not normally distributed, and non-parametric tests were conducted. We conducted a two-sample Wilcoxon Rank Sum test and Kruskal Wallis Rank test for more than 2 samples to assess the relationship between HIV prevention knowledge, and student's demographic characteristics and sexual behaviors. Spearman rank correlation test was done for continuous covariates. Library (PMCMR) in R studio software [24] was used to perform post-hoc tests using Bonferroni adjustment. Chi-square and Fisher's tests of association were also conducted to assess the association between student's behaviors and sexual behaviors. We used an iteratively reweighted least square (ILRS) generalized linear model with an identity function [25] to conduct multivariable regression analysis with knowledge score as the outcome variable. Dummy variables were generated for covariates with more than two categories. We used a 5% level of significance.

Ethical consideration

The study received ethics approval from the Joint Clinical Research Centre (JCRC) Institutional Review Board and the

Uganda National Council of Science and Technology (UNCST). Participants provided written informed consent before being interviewed. Participants were allowed to withdraw consent at any stage of the interview.

RESULTS

Socio-demographic characteristics

Socio-demographic characteristics are summarized in table 1. The mean age of the respondents was 21.2 SD (1.6) in the range of 18 to 30. More than half of the respondents 700 (52.4%) were males, and most of them 713 (53.3%) resided in university halls. Over half of the students, 801 (59.9%) were residents of urban areas, and the majority of the participants 1,208 (90.3%) reported being single. A larger proportion of our participants 1,223 (91.5%) were Christians. A total of 562 (42.0%) respondents were in the first academic year and 471 (35.2%) were in the second year. More than half of the students 946 (70.8%) were privately-sponsored, and 1,181 (88.3%) were not involved in any form of work.

HIV/AIDS prevention knowledge

The median knowledge score of students was 13.0 IQR [11-15] in the range of 0 to 18 which translates to 72.2% median pass rate. Table 2 shows the distribution of individual HIV KQ18 questions by correct scores. The table results show that HIV questions qn10 (12%), qn12 (36%), qn9 (40%), and qn17 (50%) had the lowest proportion of correct scores respectively while questions qn14 (94%), qn6 (93%), qn2 (89%) had the highest proportion of correct scores respectively.

Association between socio-demographic characteristics and HIV prevention Knowledge

Table 1 results also show the association between student's HIV prevention knowledge score and socio-demographic characteristics. The results revealed that males scored higher than females (13.0 IQR [12-15] vs. 12.0 IQR [10-14] $p=0.000$), increase in age was significantly associated with higher knowledge scores ($Rho = 0.101$, $p = 0.000$). University residence exhibited a significant relationship with HIV knowledge score ($\chi^2 = 13.2$, $p = 0.004$). Post-hoc test results showed that only scores of students residing on campus halls 13 IQR [11-15] vs. those residing at off-campus hostels 13 IQR [11-14] and parent's home 12 IQR [10-14] were significantly different from each other. The academic year of study also showed a significant association with HIV prevention knowledge ($\chi^2 = 20.1$, $p = 0.000$). The post-hoc test

TABLE 1. Socio-demographic characteristics of study population showing associations with HIV knowledge scores reporting means and proportions

Characteristic	Overall, N (%)	Knowledge score by group	
Age , mean (SD)	21.2 (1.6)	Rho=0.101*, p = 0.000	
Gender	N (%)	Median (IQR)	Statistic , p value
Male	700(52.4)	13 (12-15)	Z=6.3*, p = 0.000
Female	637(47.6)	12 (10-14)	
University residence			
Hall on campus	713(53.3)	13 (11-15)	$\chi^2 = 13.2^*$, p = 0.004
Off-campus hostel	423(31.6)	13 (11-14)	
Own home/Rental	58(4.3)	13 (11-14)	
At parent's home	143(10.7)	12 (10-14)	
Home area of residence			
Urban	801(59.9)	13 (11-15)	$\chi^2 = 3.7$, p = 0.156
Semi-urban	385(28.8)	13 (11-14)	
Rural	151(11.3)	13 (11-14)	
Marital status			
Single	1,208(90.3)	13 (11-14)	$\chi^2 = 2.9$, p = 0.402
Married	10(0.8)	14 (11-15)	
Divorced	6(0.5)	13.5 (8-15)	
Others	113(8.4)	13 (12-15)	
Year of study			
First	562(42.0)	13 (10-14)	$\chi^2 = 20.1^*$, p = 0.000
Second	471(35.2)	13 (11-14)	
Third	275(20.6)	13 (11-15)	
Fourth	29(2.2)	14 (12-15)	
Religion			
Christian	1,223(91.5)	13 (11-15)	$\chi^2 = 2.4$, p = 0.304
Muslim	95(7.1)	12 (10-14)	
Other	19(1.4)	13 (12-15)	
Sponsorship			
Private	946(70.8)	13 (11-14)	$\chi^2 = 13.6^*$, p = 0.001
Government	380(28.4)	13 (12-15)	
Other	11(0.8)	12 (10-13)	
Work			
None	1,181(88.3)	13 (11-14)	$\chi^2 = 5.4$, p = 0.066
Work part-time	150(11.2)	13 (12-15)	
Work full time	6(0.5)	15 (14-15)	

Rho -Spearman's rank correlation coefficient | IQR-Inter Quartile Range | Z- Computed Z statistic | χ^2 - Computed Chi-square statistic | * means significant at 5% level of significance

indicated that only students in the third year 13 IQR [11-15] significantly scored higher than those in the first year 13 IQR [10-14]. Student's sponsorship also posed a significant association with knowledge score ($\chi^2 = 13.6$, p = 0.001) with results from post-hoc tests indicating that government-sponsored students significantly scoring higher than privately sponsorship students. Home area of

residence, marital status, religion, and work showed no significant relationship with student's knowledge score.

We also performed an iteratively reweighted least square (ILRS) generalized linear model analysis. Table 3 shows findings from the multivariable regression analysis. The results show that after adjusting for other demographic characteristics, female students still significantly scored lower

TABLE 2. Distribution of individual HIV KQ18 questions by correct answers

HIV knowledge questions(KQ18)	Proportion	95% CI
1. Coughing and sneezing DO NOT spread HIV	0.74	0.72-0.76
2. A person can get HIV by sharing a glass of water with someone who has HIV	0.89	0.88-0.91
3. Pulling out the penis before a man climaxes/cams keeps a woman from getting HIV during sex	0.76	0.74-0.78
4. A woman can get HIV if she has anal sex with a man	0.67	0.64-0.70
5. Showering, or washing one's genitals/private parts, after sex keeps a person from getting HIV	0.89	0.87-0.90
6. All pregnant women infected with HIV will have babies born with AIDS	0.93	0.92-0.94
7. People who have been infected with HIV quickly show serious signs of being infected	0.88	0.86-0.89
8. There is a vaccine that can stop adults from getting HIV	0.74	0.71-0.76
9. People are likely to get HIV by deep kissing, putting their tongue in their partner's mouth, if their partner has HIV	0.40	0.37-0.42
10. A woman can get HIV if she has sex during her period	0.12	0.10-0.14
11. There is a female condom that can help decrease a woman's chance of getting HIV	0.832	0.81-0.85
12. A natural skin condom works better against HIV than does a latex condom	0.36	0.34-0.39
13. A person will NOT get HIV if she or he is taking antibiotics	0.80	0.78-0.82
14. Having sex with more than one partner can increase a person's chance of being infected with HIV	0.94	0.93-0.95
15. Taking a test for HIV one week after having sex will tell a person if she or he has HIV	0.53	0.50-0.55
16. A person can get HIV by sitting in a hot tub or a swimming pool with a person who has HIV	0.87	0.85-0.88
17. A person can get HIV from oral sex	0.50	0.47-0.52
18. Using Vaseline or baby oil with condoms lowers the chance of getting HIV	0.61	0.58-0.63

than males ($\beta=-0.82$, $p = 0.000$), students residing at parent's home significantly scored lower than those residing in campus halls ($\beta=-0.56$, $p=0.032$), third-year students significantly scored higher than first-year students ($\beta=0.62$, $p=0.009$) and government-sponsored students significantly scored higher than privately sponsored students ($\beta=0.54$, $p=0.002$).

Sexual and HIV preventive behaviors

Table 4 shows the distribution of sexual and HIV preventive behaviors. A total of 586 (47.9%) students reported never having at the time of the interview, and more than a half of the students 402 (55.1%) reported using a condom at the last sexual intercourse. Additionally, 55.1% of the students having had had sex with their boyfriends/girlfriends and only 8.8% reported having had a sexual encounter. A smaller proportion of respondents (6.0%) reported being drunk at their last sexual intercourse and only 21 (3.0%) students were paid in exchange for sex.

The majority of the students 1, 086 (82.5%) self-reported knowing their HIV status. More than half of the male students 420 (60.4%) reported being circumcised and 106 (44.7%) of those uncircumcised were willing to get circumcised. Results also indicate that majority of the students 1,089 (82.7%) were aware of the places where to acquire condoms and 1,037 (79.2%) students were willing to acquire condoms from those locations.

Association between student's HIV prevention knowledge, and risky sexual behaviors

Table 4 also shows the association between student's HIV knowledge and sexual behaviors. Sexual experience showed a significant relationship with HIV knowledge ($\chi^2=17.7$, $p =0.000$). Bonferroni post-hoc test results shown that scores of students who had ever had sex 13 IQR [12-15] were significantly scored higher than those who had never 12.5 IQR [10-14] and those who preferred not to answer 13 IQR [11-14]. The results also suggested that condom use experience at last sexual intercourse were significantly correlated with knowledge score ($\chi^2=15.0$, $p =0.001$). Post-hoc results showed that scores of students who preferred not to answer 12 IQR [10-14] were significantly lower than those who never used a condom 13 IQR [12-15] and those who used a condom 13 IQR [11-15]. However, no significant difference existed between those who used and never used a condom. Alcohol use at last sexual intercourse ($\chi^2=13.1$, $p =0.001$) was also significantly related knowledge score. Post-hoc results suggest that students who preferred not to answer 12 IQR [10-14] significantly scored lower than those who used 14 IQR [12-15] and never used alcohol 13 IQR [11-15]. However, there was no significant difference between those who used and never used alcohol at last sexual intercourse. Additionally, results from post-hoc test suggested there was no significant knowledge difference between those who were paid and not paid in exchange for sex.

Post-hoc tests also showed that HIV knowledge

TABLE 3. Multivariate regression analysis of association between knowledge score and demographic characteristics

Characteristic	Coefficient (SE)	95% CI
Age	-0.03 (0.06)	(-0.15,0.09)
Gender	Ref=Male	
Female	-0.82(0.16)*	(-1.14,-0.51)
University residence	Ref= on campus hall	
Off-campus hostel	0.14 (0.18)	(-0.21,0.48)
Own home/Rental	-0.31 (0.39)	(-1.07,0.44)
At parent's home	-0.56 (0.26)*	(-1.07,-0.05)
Home area of residence	Ref=Urban	
Semi-urban	-0.41 (0.17)	(-0.74,-0.07)
Rural	-0.23 (0.24)	(-0.71,0.25)
Marital status	Ref=Single	
Married	0.85 (0.92)	(-0.95,2.64)
Divorced	-0.47 (1.12)	(-2.67,1.73)
Others	0.37 (0.27)	(-1.15,0.90)
Year of study	Ref= First year	
Second	0.35 (0.18)	(-0.01, 0.70)
Third	0.62 (0.24)*	(0.16, 1.08)
Fourth	0.90 (0.56)	(-0.20, 1.99)
Religion	Ref=Christian	
Muslim	-0.24 (0.29)	(-0.82, 0.03)
Other	0.05 (0.64)	(-1.20, 1.29)
Sponsorship	Ref= Private	
Government	0.54 (0.18)*	(0.19, 0.89)
Other	-0.80 (0.83)	(-2.43, 0.83)
Work	Ref= None	
Work part-time	0.27 (0.24)	(-0.21, 0.74)
Work full time	0.94 (1.13)	(-1.27, 3.15)

* means significant at 5% level of significance

for students who were aware of places where to pick condoms 13 IQR [11-15] was significantly higher than those who weren't aware 12 IQR [9-13]. Similarly, HIV knowledge was significantly higher among students who were willing to acquire condoms 13 [11-15] than those who were not 12 IQR [10-14]. Our findings show no evidence to suggest that knowledge of HIV status, circumcision, relationship with the person at last sexual intercourse, and duration of last sexual intercourse were related to HIV knowledge score.

We further conducted an IRLS generalized linear model analysis. The results from the IRLS model are presented in Table 5. The findings show that after adjusting for other sexual behaviors showed that only knowledge of the place where to pick condoms and willingness to use condoms remained significantly associated with HIV knowledge scores. Students who knew where to pick condoms significantly scored higher than those never knew ($\beta=1.9$, $p=000$), and similarly,

students who were willing to use condoms significantly scored higher those who were unwilling ($\beta=0.9$, $p=0.029$).

Association between socio-demographic characteristics on sexual and HIV preventive behaviors

Table 6 spells out the association between socio-demographic characteristics, and sexual and HIV preventive behaviors. Older students were more likely to have ever had sex compared to younger students (31.4% vs. 17.3%), and males were more likely to have ever had sex than females (31.7% vs. 12.7%). Students staying in their own homes were more likely to have ever had sex than their counterparts. Year of study and religion were the only socio-demographic characteristics that were not significantly associated with ever having had sex. Condom use was significantly associated with age, sex, the home area of residence, year of study,

TABLE 4. Distribution of sexual and HIV preventive behaviors and their associations with HIV knowledge score

Characteristic	Overall, N(%)	Knowledge score by group	
		Median (IQR)	Statistic, p value
Ever had sex, n (%)	N=1,222		
Ever	275(22.5)	13 (12-15)	$\chi^2 = 17.7^*$, p = 0.000
Never	586 (47.9)	12.5 (10-14)	
Prefer not to answer	361(29.5)	13 (11-14)	
Number of days last had sex , median (IQR)	30 (7-120)	Rho = -0.05, p=0.423	
Condom use at last sexual intercourse, n (%)	N=730		
No	169 (23.1)	13 (12-15)	$\chi^2 = 15.0^*$, p = 0.001
Yes	402 (55.1)	13 (11-15)	
Prefer not answer	159 (21.8)	12 (10-14)	
Relationship with person of last sexual intercourse, n (%)	N=724		
Live in Partner	71 (9.8)	14 (12-15)	$\chi^2 = 10.7$, p = 0.057
Boyfriend/Girlfriend, not live in	399 (55.1)	13 (11-15)	
Casual acquaintance	64 (8.8)	13 (11.5-15)	
Prostitute	7 (1.0)	13 (11-15)	
Other	12 (1.7)	13.5 (12.5-15)	
Prefer not to answer	171 (23.6)	13 (10-15)	
Alcohol use at last sexual intercourse, n (%)	N=710		
No	553 (77.9)	13 (11-15)	$\chi^2 = 13.1^*$, p = 0.001
Yes	43 (6.1)	14 (12-15)	
Prefer not to answer	114 (16.1)	12 (10-14)	
If yes, who was drunk, n (%)	N=49		
Respondent	8 (16.3)	14 (13.5-14.5)	$\chi^2 = 2.9$, p = 0.414
Partner	11 (22.4)	15 (12-16)	
Both	15 (30.6)	14 (10-15)	
Prefer not to answer	15(30.6)	13 (10-14)	
Pay (got paid) in sex exchange, n (%)	N=698		
No	588 (84.2)	13 (11-15)	$\chi^2 = 23.5^*$, p = 0.000
Yes	21 (3.0)	12 (10-15)	
Prefer not to answer	89 (12.8)	12 (9-14)	
Knowledge of HIV status, n (%)	N=1,316		
No	177 (13.5)	13 (11-14)	$\chi^2 = 6.2$, p = 0.050
Yes	1,086 (82.5)	13 (11-15)	
Prefer not to answer	53 (4.0)	12 (10-14)	
Circumcision [Males], n (%)	N=695		
No	237 (34.1)	13 (12-15)	$\chi^2 = 1.8$, p = 0.403
Yes	420 (60.4)	13 (11-15)	
Prefer not to answer	38 (5.5)	12.5 (11-15)	
Willingness to get circumcised, n (%)	N=237		
No	117 (48.9)	13 (11-15)	$\chi^2 = 0.7$, p = 0.711
Yes	107 (44.8)	14 (11-14)	
Prefer not to answer	15 (6.3)	13 (11-14)	
Knowledge of places where to pick condoms, n (%)	N=1,317		
No	118 (9.0)	12 (9-13)	$\chi^2 = 43.1^*$, p = 0.000
Yes	1,089 (82.7)	13 (11-15)	
Prefer not to answer	110 (8.3)	11.5 (9-14)	
Willingness to acquire a condom, n (%)	N=1,309		
No	150 (11.5)	12 (10-14)	$\chi^2 = 41.5^*$, p = 0.000
Yes	1,037 (79.2)	13 (11-15)	
Prefer not to answer	122 (9.3)	11 (9-14)	

* means significant at 5% level of significance | IQR-Inter Quartile Range | χ^2 -Chi-square value

TABLE 5. Multivariate regression analysis of association between knowledge score and HIV prevention and sexual behaviors

Characteristic	Coefficient (SE)	95% CI
Ever had sexual intercourse	Ref=Ever	
Never	0.05 (0.25)	(-0.45,0.54)
Prefer not answer	omitted	
Condom use at last sexual intercourse	Ref= Not used	
Yes	-0.40 (0.28)	(-0.95,0.15)
Prefer not answer	-0.54 (0.43)	(-1.38,0.30)
Alcohol use at last sexual intercourse	Ref=No	
Yes	0.19 (0.48)	(-0.75, 1.14)
Prefer not answer	0.11 (0.45)	(-0.81,1.04)
Pay (got paid) in sex exchange	Ref=No	
Yes	-1.28 (0.68)	(-2.62,0.06)
Prefer not answer	-1.42* (0.45)	(-2.31,-0.54)
Knowledge of places where to pick condoms	Ref= No	
Yes	1.90* (0.44)	(1.02,2.77)
Prefer not to answer	1.74* (0.71)	(0.35,3.13)
Willingness to acquire a condom	Ref=No	
Yes	0.91* (0.42)	(0.10,1.73)
Prefer not to answer	-0.16 (0.65)	(-1.43,1.11)

Other sexual and prevention variables were excluded due to collinearity

* means significant at 5% level of significance

student’s sponsorship and work. More males reported condom use compared to females (63.3% vs. 55.1%), and students in the third year of study were more likely to use condoms compared to those in other academic years. Knowledge of HIV status showed no significant association with the socio-demographic characteristics. Circumcision was significantly associated with marital status, year of study, religion, work and home area of residence. Students from urban centers and those not in relationships were more likely to be circumcised than their counterparts. Willingness to get circumcised was only significantly related to student’s sponsorship type and home area of residence.

DISCUSSION

In this study, we assessed the predictors of HIV prevention knowledge and sexual behaviors among students aged 18 to 30 years at Makerere University Kampala, Uganda.

Our findings showed that median knowledge score was 13.0 IQR (11-15) on a scale of 0 to 18, an indication that averagely students scored about 72%. This reveals that Makerere University student’s HIV knowledge was relatively good. This finding correlates very well with a study conducted in Belgium that reported a fair knowledge score of 20 IQR [19-21] among university students [26], and

other similar studies elsewhere [27,28], but it differs with prior reviews in Ethiopia [29] and Sudan [30] that reported a generally poor performance among university students.

Our study also shows that students performed poorly on HIV transmission questions, and this result is consistent with a prior study in Ethiopia [29] that revealed low knowledge on HIV transmission among adolescents. This finding is, however, inconsistent with results of a related study that conducted among high school students in Belgium and showed a higher knowledge score on HIV transmission questions [31]. Low HIV transmission knowledge scores suggest that Makerere University students are at a high risk of contracting HIV. Our study recommends the implementation of student-tailored HIV transmission educational programs in order to close this knowledge gap.

Males demonstrated a higher level of HIV knowledge than females both at univariate and multivariable analysis. This result is inconsistent with studies conducted in Iran, Belgium [31,32], and Ghana [33] that showed that females scored higher than males. Conversely, a study conducted in Ethiopia [29] revealed no gender differences in HIV Knowledge levels. Different socio-cultural norms in different countries and parts of the countries may explain these gender differences. Behavioral and educational programs that consider the gender dimensions among university students could help in closing this gap. Our study also shows that student’s HIV knowledge increased

with increase in age. This result is supported by prior studies [26,31] that suggest that older students scored higher than younger students. Our study evidenced that students residing within university housings scored better than those that resided at their parent's homes. This shows that students residing on campus may be benefiting from the various HIV social campaigns that run within the confines of the university residences. Conversely, those residing outside the university's residence miss out on such campaigns which are impacting on their low level of knowledge. Additionally, one can also argue that parents are not taking a key role in discussing HIV related information with their children. Future research could exploit ways of involving parents in championing HIV campaigns among their children. Our study also revealed that government-sponsored students scored higher than the privately sponsored student. This result can be related to that of residing in university residences. Most government-sponsored students reside in university halls and tap knowledge from the various HIV campaigns as compared to private students whom majority reside outside the university. Our study didn't find any association between knowledge and marital status. We had anticipated students in marriage to be more knowledgeable than the unmarried as reported by a prior study [26].

Over half of the respondents studied had never had sex. In line with a prior study carried out in China [34], we found evidence to suggest that most of the students reported using a condom at their last sexual debut. Very few university students were engaged in sex for money. On a positive side also, alcohol use during sex was found to be very low and Knowledge of self-reported HIV status was very high among college students. Our study also suggested that more than half of our male respondents were circumcised. Circumcision reduces the risk of HIV infection by about 60% and the national prevalence rate of circumcision stands at 26%. Males from urban areas were more likely to be circumcised compared to those from rural areas as supported by a previous study in Uganda [35]. Although self-reported circumcision was high, willingness to circumcise among the uncircumcised males was below the average, contrary to earlier studies in Rwanda [36] and Uganda [35,37].

Sexual experience was significantly related to HIV knowledge scores. Students that reported prior sexual debut scored higher than those that reported having never had any sexual encounter. This seemingly paradoxical result suggests that knowledge does not necessarily translate to safer sex practices. Similar findings have also been reported among university students in Belgium [26]. This can be attributed to personal experiences where for example sexually active students tend to seek more information on the spread and control of HIV than the non-sexually active students. This shows a need for more HIV interventions tailored to both sexually experienced and non-sexually experienced students. Our study did not find any evidence to suggest that condom use at last sexual

intercourse and knowledge of HIV status were related to HIV knowledge scores. Knowledge of places where to acquire condoms, and willingness to use condoms were associated with better HIV knowledge outcomes. This indicates that positive attitudes towards condom use are positively correlated with high HIV knowledge gains.

Our study suggests that increase in age is associated with sexual experience among university students. This result is supported by a prior study in Uganda [20]. This finding is rather not surprising, as African communities dictate that as you age, you're expected to marry, and implicit in this is sexual intercourse. Male students and those staying in their own homes were more likely to have ever had sex compared to the rest of the students. In contrast to a similar study in Uganda [19], we didn't find evidence of an association between sexual experience and religion. This difference may be due to the fact that this study never disaggregated Christianity to capture different dimensions as compared to a previous study in Uganda [19] that used various dimensions. Our study suggests that males are more likely to use condoms than females as opposed to a prior review in Tanzania [38] that indicated that females were more likely to use condoms than males. Condom awareness efforts targeting females are needed in universities in order to make them first consumers of condom use as this will trigger high safe sexual practices among university students.

Research implications for intervention and policy

The present study has demonstrated the existence of gender differences in HIV prevention knowledge and sexual behaviors among university students. Designing of innovative ways that address these gender differences across similar study groups would be fundamental in achieving safe sexual behaviors and improving HIV knowledge. Our study also shows that students residing in university housings performed better than those that reside at parent's homes. Policy should identify possible ways of involving parents in improving HIV knowledge among adolescents. Our study has shown evidence that students from urban areas were more circumcised. Similarly, willingness to circumcise was higher among students residing from urban areas compared to rural areas. Policy makers need to come up with programs that target rural population aiming at creating demand for affordable and safe male circumcision.

Study strengths and limitations

We used an internationally-validated questionnaire to measure HIV knowledge among students. This study can potentially allow cross-country comparisons of knowledge scores in similar study settings. The study questionnaires were completed by the respondents which reduced the interviewer bias that sometimes arises as a result of wrong

interpretations of questions by interviewers.

A limitation of the study is that we excluded participants who did not possess an Android smartphone and those in the final semester of their academic year. As a result, the findings may not be generalizable to all students in Makerere or students in other universities in Uganda.

CONCLUSION

Our study findings show that Makerere University students possess a good level of knowledge on HIV prevention. The study has shown evidence of associations between HIV knowledge, and sexual experience and willingness to use condoms. Our study demonstrates the existence of significant gendered HIV knowledge and sexual behavior differences among university students. Behavioral and educational Programs that target both sexually and non-sexually experienced students should address the gender differences. Future research should aim to explore ways of involving parents in improving HIV knowledge among adolescents. Future research should also aim at assessing the HIV/AIDS knowledge levels among students that use drugs, and those with other sexual orientations.

Acknowledgement

We would like to thank Grand Challenges Canada for funding this study. We would also like to recognize the contribution of our research assistants and Makerere University students that participated in this study.

Author's contributions

AN1, PN, EN, AN3, SJL, JBB conceived the study. AN1, PN, EN carried out the data collection. AN1 carried out data analysis and drafted the first version of the manuscript. AN1, ONO, PN, EN, AN3, SJL, JB edited and approved the manuscript.

Competing interest

The authors declare no competing interests.

References

1. Ministry of Health Kampala Uganda. Ministry of Health [Uganda], ICF International, CDC, USAID, WHO: Uganda AIDS Indicator Survey. Maryland: Ministry of Health and ICF International; 2012. Available from: http://health.go.ug/docs/UAIS_2011_REPORT.pdf. Available from: http://health.go.ug/docs/UAIS_2011_REPORT.pdf
2. MoH. Ministry of Health [Uganda], ORC Macro: Uganda HIV/AIDS Sero-behavioural Survey 2004-2005. Calverton, Maryland, USA: Ministry of Health and ORC Macro; 2006. Available from: <http://dhsprogram.com/pubs/pdf/AIS2/AIS2.pdf>. Available from: http://pdf.usaid.gov/pdf_docs/Pnadv508.pdf
3. UNAIDS. UNAIDS Prevention Gap Report 2016; Available from: file:///C:/Users/GHE/Desktop/KQ18/2016-prevention-gap-report_en.pdf
4. UNAIDS. GLOBAL REPORT: UNAIDS report on the global AIDS epidemic 2013 [Internet]. Unids. 2013. 198 p. Available from: www.unaids.org/.../unaids/.../2013/gr2013/UNAIDS_Global_Report_2013
5. Uganda AIDS Commission: Global AIDS Response Progress Report; 2012. Available from: [http://www.unaids.org/en/dataanalysis/knowyourresponse/countryprogressreports/2012countries/ce_UG_Narrative_Report\[1\].pdf](http://www.unaids.org/en/dataanalysis/knowyourresponse/countryprogressreports/2012countries/ce_UG_Narrative_Report[1].pdf); Accessed September 22, 2014.
6. Li S, Huang H, Xu G, Cai Y, Huang F, Ye X. Substance use, risky sexual behaviors, and their associations in a Chinese sample of senior high school students. *BMC public health*. 2013 Apr 4;13(1):295. Available from: <https://bmcpubhealth.biomedcentral.com/articles/10.1186/1471-2458-13-295>
7. Connell CM, Gilreath TD, Hansen NB. A multiprocess latent class analysis of the co-occurrence of substance use and sexual risk behavior among adolescents. *Journal of Studies on Alcohol and Drugs*. 2009 Nov;70(6):943-51. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2776124/>
8. Abebe G, Fekadu A. A health concerns and challenges among high school adolescents. *Ethiop J Health Dev*. 2000;10(1):37-40.
9. Shiferaw Y, Alemu A, Girma A, et al. Assessment of knowledge, attitude and risk behaviors towards HIV/AIDS and other sexual transmitted infection among preparatory students of Gondar town, north west Ethiopia. *BMC research notes*. 2011 Nov 21;4(1):505. Available from: <https://bmcrsnotes.biomedcentral.com/articles/10.1186/1756-0500-4-505>
10. Zinabu A. Knowledge, attitude and behavior on HIV/AIDS/STD among workers in the informal sectors in Addis Ababa. *Ethiop J Health Dev*. 2003.
11. Prince A, Bernard AL. Sexual behaviors and safer sex practices of college students on a commuter campus. *Journal of American College Health*. 1998 Jul 1;47(1):11-21.
12. Opio A, Mishra V, Hong R, et al. Trends in HIV-related behaviors and knowledge in Uganda, 1989-2005: evidence of a shift toward more risk-taking behaviors. *JAIDS Journal of Acquired Immune Deficiency Syndromes*. 2008 Nov 1;49(3):320-6.
13. Maswanya ES, Maji K, Horiguchi I, et al. Knowledge, risk perception of AIDS and reported sexual behaviour among students in secondary schools and colleges in Tanzania. *Health education research*. 1999 Apr 1;14(2):185-96. Available from: <https://academic.oup.com/her/article/14/2/185/790308>
14. Odu OO, Asekun-Olarinmoye EO, Bamidele JO, Egbewale BE, Amusan OA, Olowu AO. Knowledge, attitudes to HIV/AIDS and sexual behaviour of students in a tertiary institution in south-western Nigeria. *The European journal of contraception & reproductive health care*. 2008 Jan 1;13(1):90-6. Available from: <http://www.tandfonline.com/doi/abs/10.1080/13625180701617670?journalCode=iejc20>
15. Ntata PR, Muula AS, Siziya S, Kayambazinthu EE. Gender differences in university students' HIV/AIDS-related knowledge

- and sexual behaviours in Malawi: a pilot study: brief report. SAHARA: Journal of Social Aspects of HIV/AIDS Research Alliance. 2008 Dec 1;5(4):201-5. Available from: http://journals.co.za/docserver/fulltext/m_sahara/5/4/m_sahara_v5_n4_a5.pdf?expires=1500234393&id=id&accname=guest&checksum=F8C942C3B118C29135561F35995D66C4
16. Lema LA, Katapa RS, Musa AS. Knowledge on HIV/AIDS and sexual behaviour among youths in Kibaha District, Tanzania. *Tanzania journal of health research*. 2008;10(2):79-83. Available from:
 17. Takyi BK. Correlates of HIV/AIDS-related knowledge and preventive behavior of men in Africa. *Journal of health and human services administration*. 2001 Oct 1:234-57.
 18. Lule GS, Gruer LD. Sexual behaviour and use of the condom among Ugandan students. *AIDS care*. 1991 Jan 1;3(1):11-9.
 19. Agardh A, Tumwine G, Östergren PO. The impact of socio-demographic and religious factors upon sexual behavior among Ugandan university students. *PLoS One*. 2011 Aug 24;6(8):e23670. Available from: <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0023670>
 20. Agardh A, Cantor-Graae E, Östergren PO. Youth, sexual risk-taking behavior, and mental health: a study of university students in Uganda. *International journal of behavioral medicine*. 2012 Jun 1;19(2):208-16. Available from: <https://link.springer.com/article/10.1007/s12529-011-9159-4>
 21. Choudhry V, Agardh A, Stafström M, Östergren PO. Patterns of alcohol consumption and risky sexual behavior: a cross-sectional study among Ugandan university students. *BMC public health*. 2014 Feb 6;14(1):128. Available from: <https://bmcpublichealth.biomedcentral.com/articles/10.1186/1471-2458-14-128>
 22. Agardh A, Odberg-Pettersson K, Östergren PO. Experience of sexual coercion and risky sexual behavior among Ugandan university students. *BMC public health*. 2011 Jul 4;11(11):527. Available from: <https://bmcpublichealth.biomedcentral.com/articles/10.1186/1471-2458-11-527>
 23. Carey MP, Schroder KE. Development and psychometric evaluation of the brief HIV Knowledge Questionnaire. *AIDS education and prevention*. 2002 Apr 2;14(2):172-82. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2423729/>
 24. Pohlett T. The pairwise multiple comparison of mean ranks package (PMCMR). R package. 2014 Jun 24:2004-6. Available from: <http://cran.uni-muenster.de/web/packages/PMCMR/vignettes/PMCMR.pdf>
 25. Green PJ. Iteratively reweighted least squares for maximum likelihood estimation, and some robust and resistant alternatives. *Journal of the Royal Statistical Society. Series B (Methodological)*. 1984 Jan 1:149-92. Available from: <http://128.105.7.14/courses/st771-newton/papers/green.pdf>
 26. Degroote S, Vogelaers D, Liefhooghe G, Vermeir P, Vandijck DM. Sexual experience and HIV-related knowledge among Belgian university students: a questionnaire study. *BMC research notes*. 2014 May 15;7(1):299. Available from: <https://bmcresnotes.biomedcentral.com/articles/10.1186/1756-0500-7-299>
 27. Platten M, Pham HN, Nguyen HV. Knowledge of HIV and factors associated with attitudes towards HIV among final-year medical students at Hanoi medical university in Vietnam. *BMC public health*. 2014 Mar 20;14(1):265. Available from: <https://bmcpublichealth.biomedcentral.com/articles/10.1186/1471-2458-14-265>
 28. Nubed CK, Akoachere JF. Knowledge, attitudes and practices regarding HIV/AIDS among senior secondary school students in Fako Division, South West Region, Cameroon. *BMC public health*. 2016 Aug 22;16(1):847. Available from: https://bmcpublichealth.biomedcentral.com/articles/10.1186/s12889-016-3516-9?utm_campaign=BMC_TrendMD&utm_medium=cpc&utm_source=TrendMD
 29. Paul M, Majumdar B, Akhtar-Danesh N, Boblin S, Sherifali D. The gendered knowledge of HIV/AIDS among urban University students in Southern Ethiopia. *European Scientific Journal, ESJ*. 2013 Jul 12;9(21).
 30. Nasir EF, Åström AN, David J, Ali RW. HIV and AIDS related knowledge, sources of information, and reported need for further education among dental students in Sudan-a cross sectional study. *BMC Public Health*. 2008 Aug 14;8(1):286. Available from: <https://bmcpublichealth.biomedcentral.com/articles/10.1186/1471-2458-8-286>
 31. Van Rossem R, Berten H, Van Tuyckom C. AIDS knowledge and sexual activity among Flemish secondary school students: a multilevel analysis of the effects of type of education. *BMC public health*. 2010 Jan 21;10(1):30. Available from: <https://bmcpublichealth.biomedcentral.com/articles/10.1186/1471-2458-10-30>
 32. Tavoosi A, Zaferani A, Enzevaei A, Tajik P, Ahmadinezhad Z. Knowledge and attitude towards HIV/AIDS among Iranian students. *BMC public health*. 2004 May 24;4(1):17. Available from: <https://bmcpublichealth.biomedcentral.com/articles/10.1186/1471-2458-4-17>
 33. Oppong AK, Oti-Boadi M. HIV/AIDS knowledge among undergraduate university students: implications for health education programs in Ghana. *African health sciences*. 2013;13(2):270-7. Available from: <https://www.ajol.info/index.php/ahs/article/viewFile/93478/82904>
 34. Long L, Yuan T, Wang M, et al. Factors associated with condom use among male college students in Wuhan, China. *PLoS one*. 2012 Dec 18;7(12):e51782. Available from: <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0051782>
 35. Wilcken A, Miiro-Nakayima F, Hizaamu RN, Keil T, Balaba-Biyansi D. Male circumcision for HIV prevention-a cross-sectional study on awareness among young people and adults in rural Uganda. *BMC public health*. 2010 Apr 26;10(1):209. Available from: <https://bmcpublichealth.biomedcentral.com/articles/10.1186/1471-2458-10-209>
 36. Gasasira RA, Sarker M, Tsague L, et al. Determinants of circumcision and willingness to be circumcised by Rwandan men, 2010. *BMC public health*. 2012 Feb 18;12(1):134. Available from: <https://bmcpublichealth.biomedcentral.com/articles/10.1186/1471-2458-12-134>
 37. Kibira SP, Makumbi F, Daniel M, Atuyambe LM, Sandøy IF. Sexual Risk Behaviours and Willingness to Be Circumcised among Uncircumcised Adult Men in Uganda. *PLoS one*. 2015 Dec 14;10(12):e0144843. Available from: <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0144843>
 38. Namisi FS, Aarø LE, Kaaya S, Onya HE, Wubs A, Mathews C. Condom use and sexuality communication with adults: a study among high school students in South Africa and Tanzania. *BMC public health*. 2013 Sep 23;13(1):874. Available from: <https://bmcpublichealth.biomedcentral.com/articles/10.1186/1471-2458-13-874>

SUPPLEMENTARY FILES

1. Table

TABLE 6. Association between socio-demographic characteristics and sexual and HIV preventive behaviors

Characteristic	Ever had sex			Condom use at last sexual intercourse			Knowledge of HIV status			Circumcision [Males only]		
	Ever	Never	Prefer not to answer	No	Yes	Prefer not to answer	No	Yes	Prefer not to answer	No	Yes	Prefer not to answer
Age												
18-21	133(17.3)	427(55.4)	210(27.3)	73(19.0)	210(54.5)	102(26.5)	119(14.6)	665(81.7)	30(3.7)	116(32.3)	205(60.6)	17(5.0)
22-30	142(31.4)	159(35.2)	151(33.4)	96(27.1)	192(55.6)	57(21.9)	58(11.5)	421(83.9)	23(4.6)	121(33.9)	215(60.2)	21(5.9)
	P value= 0.000**			P value = 0.001**			P value = 0.227			P value = 0.885		
Gender												
Male	200(31.7)	212(33.6)	219(34.7)	108(22.6)	302(63.3)	67(14.1)	95(13.8)	565(82.0)	29(4.2)	237(34.1)	420(60.4)	38(5.8)
Female	75(12.7)	374(63.3)	142(29.5)	61(24.1)	402(55.1)	92(36.4)	82(13.1)	521(83.1)	24(4.0)			
	P value= 0.000**			P value = 0.000**			P value = 0.866					
University residence												
Hall on campus	140(21.2)	338(51.2)	181(27.5)	78(21.5)	205(56.5)	80(22.0)	94(13.4)	582(82.9)	26(3.7)	128(33.6)	231(60.6)	22(5.8)
Off-campus hostel	97(25.3)	164(42.8)	122(31.8)	62(24.6)	139(55.2)	51(20.2)	53(12.7)	348(83.6)	15(3.6)	79(34.8)	137(60.3)	11(4.8)
Own home/Rental	19(38.0)	15(30.0)	16(32.0)	11(25.6)	27(62.8)	5(11.6)	11(19.3)	43(75.4)	3(5.3)	15(34.1)	26(59.1)	3(6.8)
At parent's home	19(14.6)	69(53.1)	42(32.3)	18(25.0)	31(43.1)	23(31.9)	19(13.5)	113(80.1)	9(6.4)	15(34.9)	26(60.5)	2(4.6)
	P value= 0.002**			P value = 0.168			P value = 0.589			P value = 0.998		
Home area of residence												
Urban	147(20.1)	366(49.9)	220(30.0)	91(21.6)	234(55.6)	96(22.8)	102(13.0)	649(82.7)	34(4.3)	133(33.6)	248(62.6)	15(3.8)
Semi-urban	82(23.7)	153(44.2)	111(32.1)	47(20.5)	133(58.1)	49(21.4)	52(13.6)	316(82.7)	14(3.7)	63(31.0)	121(59.6)	19(9.4)
Rural	46(32.2)	67(46.8)	30(21.0)	31(38.7)	35(43.7)	14(17.5)	23(15.4)	121(81.2)	5(3.4)	41(42.7)	51(53.1)	4(4.2)
	P value = 0.007**			P value = 0.013*			P value = 0.899			P value = 0.019*		
Marital status												
Single	240(21.6)	551(49.7)	318(28.7)	145(22.7)	354(55.4)	140(21.9)	161(13.5)	984(82.8)	44(3.7)	222(34.8)	385(60.4)	30(4.7)
Married	4(50.0)	1(12.5)	3(37.5)	6(66.7)	3(33.3)	0(0.0)	1(10.0)	8(80.0)	1(100.0)	3(42.9)	3(42.7)	1(14.3)
Divorced	0(0.0)	2(33.3)	4(66.7)	1(25.0)	1(25.0)	2(50.0)	0(0.0)	6(100.0)	0(0.0)	0(0.0)	4(100.0)	0(0.0)
Others	31(31.3)	32(32.3)	36(36.4)	17(21.8)	44(56.4)	17(21.8)	15(13.5)	88(79.3)	8(7.2)	12(25.5)	28(59.6)	7(14.9)
	P value = 0.002**			P value = 0.054			P value = 0.480			P value = 0.035*		
Year of study												
First	111(21.4)	274(52.9)	133(31.8)	50(18.0)	156(56.1)	72(25.9)	77(14.0)	456(83.2)	15(2.7)	94(34.7)	173(63.8)	4(1.5)
Second	94(22.2)	195(46.0)	135(31.8)	76(27.8)	144(52.7)	53(19.4)	62(13.2)	384(81.7)	24(5.1)	82(36.0)	128(56.1)	18(7.9)
Third	64(25.2)	106(41.7)	84(33.1)	35(21.3)	98(59.8)	31(18.9)	35(12.9)	222(81.9)	14(5.2)	55(31.6)	105(60.3)	14(8.1)
Fourth	6(23.1)	11(42.3)	9(34.6)	8(53.3)	4(26.7)	3(20.0)	3(11.1)	24(88.9)	0(0.0)	6(27.3)	14(60.4)	2(9.1)
	P value = 0.089			P value = 0.005**			P value = 0.408			P value = 0.019*		
Religion												
Christian	253(22.7)	539(48.3)	324(29.0)	152(22.9)	365(54.9)	148(22.3)	158(13.1)	995(82.7)	50(4.2)	233(36.2)	374(58.2)	36(5.6)
Muslim	20(22.5)	39(43.8)	30(33.7)	14(25.5)	33(60.0)	8(14.5)	16(17.1)	76(80.8)	2(2.1)	0(0.0)	38(61.5)	1(2.6)
Other	2(11.8)	8(47.1)	7(41.2)	3(30.0)	4(40.0)	3(30.0)	3(15.8)	15(78.9)	1(5.3)	4(30.8)	8(61.5)	1(7.7)
	P value = 0.621			P value = 0.607			P value = 0.719			P value = 0.000**		
Sponsorship												
Private	203(23.7)	382(44.7)	270(31.6)	122(22.1)	314(56.9)	116(21.0)	128(13.7)	765(82.2)	38(4.1)	161(33.7)	292(61.1)	25(5.2)
Government	70(19.6)	196(54.9)	91(25.5)	45(25.6)	88(50.0)	43(24.4)	46(12.2)	315(83.8)	15(4.0)	72(34.1)	126(59.7)	13(6.2)
Other	2(20.0)	8(80.0)	0(0.0)	2(100.0)	0(0.0)	0(0.0)	3(33.3)	6(66.7)	0(0.0)	4(66.7)	2(33.3)	0(0.0)
	P value = 0.003**			P value = 0.056			P value = 0.427			P value = 0.519		
Work												
None	235(21.7)	535(49.4)	312(28.8)	133(21.3)	354(56.5)	139(22.2)	159(13.7)	957(82.3)	47(4.0)	215(36.3)	345(58.2)	33(5.6)
Work parttime	40(29.6)	49(36.3)	46(34.1)	34(34.0)	46(46.0)	20(20.0)	18(12.2)	123(83.7)	6(4.1)	22(22.5)	71(72.4)	5(5.1)
Work full time	0(0.0)	2(40.0)	3(60.0)	2(50.0)	2(50.0)	0(0.0)	0(0.0)	6(100.0)	0(0.0)	0(0.0)	4(100.0)	0(0.0)
	P value = 0.021*			P value = 0.038*			P value = 0.826			P value = 0.037*		

** means significant at 1% level of significance and * means significant at 5% level of significance

TABLE 6 (Cont'd). Association between socio-demographic characteristics and sexual and HIV preventive behaviors

Characteristic	Willingness to circumcise (Males only)			Alcohol use at last sexual intercourse			Paid in exchange for sex			Knowledge of places where to pick condoms		
	No	Yes	Prefer not to answer	No	Yes	Prefer not to answer	No	Yes	Prefer not to answer	No	Yes	Prefer not to answer
Age												
18-21	52(44.4)	57(48.7)	8(6.9)	279(74.6)	21(5.6)	74(19.8)	304(82.6)	10(2.7)	54(14.7)	69(8.5)	669(82.3)	75(9.2)
22-30	65(53.3)	50(41.0)	7(5.7)	274(81.5)	22(6.6)	40(11.9)	284(86.1)	11(3.3)	35(10.6)	49(9.7)	420(83.3)	35(6.9)
	P value = 0.394			P value = 0.017*			P value = 0.256			P value = 0.285		
Gender												
Male	117(49.0)	107(44.8)	15(6.3)	384(82.9)	32(6.9)	47(10.2)	400(87.3)	17(3.7)	41(9.0)	55(7.9)	611(87.7)	31(4.5)
Female	[Males only]			169(68.4)	11(4.5)	67(27.1)	188(78.3)	4(1.7)	48(20.0)	63(10.2)	478(77.1)	79(12.7)
				P value = 0.000**			P value = 0.000**			P value = 0.000**		
University residence												
Hall on campus	66(51.2)	57(44.2)	6(4.6)	275(78.1)	18(5.1)	59(16.8)	289(84.3)	11(3.2)	43(12.5)	49(6.9)	596(84.4)	61(8.6)
Off-campus hostel	37(47.4)	37(47.4)	4(5.1)	189(77.1)	21(8.6)	35(14.3)	205(83.7)	10(4.1)	30(12.2)	47(11.3)	340(82.3)	26(6.3)
Own home/Rental	9(56.3)	4(25.0)	3(18.7)	36(85.7)	2(4.8)	4(9.5)	37(92.5)	0(0.0)	3(7.5)	6(10.3)	47(81.0)	5(8.6)
At parent's home	5(31.3)	9(56.2)	2(12.5)	53(74.6)	2(2.8)	16(22.5)	57(81.4)	0(0.0)	13(18.6)	16(11.4)	106(75.7)	18(12.9)
	P value = 0.172			P value = 0.208			P value = 0.279			P value = 0.034*		
Home area of residence												
Urban	57(42.9)	70(52.6)	6(4.5)	323(79.0)	24(5.9)	62(15.2)	331(82.7)	13(3.3)	56(14.0)	68(8.6)	660(83.3)	65(8.2)
Semi-urban	32(49.2)	26(40.0)	7(10.8)	166(74.8)	13(5.9)	43(19.4)	188(85.8)	5(2.3)	26(11.9)	35(9.4)	307(82.1)	32(8.6)
Rural	28(68.3)	11(26.8)	2(4.9)	64(81.0)	6(7.6)	9(11.4)	69(87.3)	3(3.8)	7(8.9)	15(10.0)	122(81.3)	13(8.7)
	P value = 0.016*			P value = 0.466			P value = 0.654			P value = 0.972		
Marital status												
Single	108(48.6)	100(45.1)	14(6.3)	486(78.3)	36(5.8)	99(15.9)	518(84.9)	19(3.1)	73(12.0)	108(9.1)	986(82.6)	97(8.1)
Married	2(66.7)	1(33.3)	0(0.0)	7(77.8)	2(22.2)	0(0.0)	6(66.7)	2(22.2)	1(11.1)	0(0.0)	9(90.0)	1(10.0)
Divorced	0(0.0)	0(0.0)	0(0.0)	2(50.0)	2(50.0)	0(0.0)	2(50.0)	0(0.0)	2(50.0)	1(16.7)	4(66.7)	1(16.7)
Others	7(50.0)	6(42.9)	1(7.1)	58(76.3)	3(3.9)	15(19.7)	62(82.7)	0(0.0)	13(17.3)	9(8.2)	90(81.2)	11(10.0)
	P value = 0.973			P value = 0.002**			P value = 0.002**			P value = 0.858		
Year of study												
First	47(49.5)	44(46.3)	4(4.2)	207(76.1)	17(6.2)	48(17.6)	217(82.0)	9(3.4)	38(14.4)	56(10.2)	446(80.9)	49(8.9)
Second	38(45.2)	40(47.6)	6(7.1)	206(78.0)	15(5.7)	43(16.3)	220(84.0)	7(2.7)	35(13.4)	42(9.0)	386(82.8)	38(8.2)
Third	30(55.6)	20(37.0)	4(7.4)	128(80.0)	10(6.3)	22(13.7)	138(87.3)	5(3.2)	15(9.5)	18(6.6)	232(85.6)	21(7.8)
Fourth	2(33.3)	3(50.0)	1(16.7)	12(86.7)	1(7.1)	1(7.1)	13(92.9)	0(0.0)	1(7.1)	2(6.9)	25(86.2)	2(6.9)
	P value = 0.694			P value = 0.911			P value = 0.757			P value = 0.732		
Religion												
Christian	114(48.5)	107(45.5)	14(6.0)	507(77.9)	37(5.7)	107(16.4)	534(84.4)	18(2.8)	81(12.8)	105(8.7)	1,004(83.3)	97(8.0)
Muslim	0(0.0)	0(0.0)	1(25.0)	41(80.4)	4(7.8)	6(11.8)	49(89.1)	1(1.8)	5(9.1)	10(10.9)	74(80.4)	8(8.7)
Other	75(75.0)	0(0.0)	1(25.0)	5(62.5)	2(25.0)	1(12.5)	5(50.0)	2(20.0)	3(30.0)	3(15.8)	11(57.9)	5(26.3)
	P value = 0.097			P value = 0.188			P value = 0.006**			P value = 0.033*		
Sponsorship												
Private	79(48.5)	70(42.9)	14(8.6)	423(78.8)	33(6.2)	81(15.1)	448(84.5)	15(2.8)	67(12.6)	84(9.0)	773(83.2)	72(7.8)
Government	35(48.6)	37(51.4)	0(0.0)	129(75.0)	10(5.8)	33(19.2)	139(83.7)	5(3.0)	22(13.3)	29(7.7)	311(82.5)	37(9.8)
Other	3(75.0)	0(0.0)	1(25.0)	1(100.0)	0(0.0)	0(0.0)	1(50.0)	1(50.0)	0(0.0)	5(7.7)	5(45.5)	1(9.1)
	P value = 0.023*			P value = 0.752			P value = 0.004**			P value = 0.000**		
Work												
None	106(48.8)	97(44.7)	14(6.5)	472(77.6)	34(5.6)	102(16.8)	507(84.6)	19(3.2)	73(12.2)	111(9.6)	955(82.2)	96(8.3)
Work part-time	11(50.0)	10(45.5)	1(4.5)	78(79.6)	8(8.2)	12(12.2)	78(82.1)	1(1.1)	16(16.8)	7(4.7)	128(85.9)	14(9.4)
Work full time	0(0.0)	0(0.0)	0(0.0)	3(75.0)	1(25.0)	0(0.0)	3(75.0)	1(25.0)	0(0.0)	0(0.0)	6(100.0)	0(0.0)
	P value = 0.940			P value = 0.280			P value = 0.046*			P value = 0.071		

** means significant at 1% level of significance and * means significant at 5% level of significance

2. Research instrument

Development and Pilot Testing of a Mobile Phone Gaming Application to Increase Knowledge of HIV Prevention among Students at Makerere University, Kampala

Date: |__ __|__ __|__ __ __ __|

Participant #: |__ __ __|

Cell Phone #: |_____|

SOCIO-DEMOGRAPHICS

1. Age: |__ __| years
2. Gender: Male |__| Female |__|
3. Residence: Hall on campus |__| Off-campus hostel |__| own home |__| At parents' |__|
4. Original place of residence: Urban |__| Semi-urban |__| Rural |__|
5. Marital status: Single |__| Married |__| Divorced |__|
6. Year: First |__| Second |__| Third |__| Fourth |__| Fifth |__|
7. Religion: Christian |__| Muslim |__| Other |__|
8. Type of sponsorship: Private |__| Government |__| Other |__|
9. Any work? None |__| Work part-time |__| Work full time |__|
10. If you are a paid employee, how much do you earn per month? |__ __ __ __ __ __ __ __|
11. Asset ownership (tick all that apply): Land |__| House |__| Car |__| Motorcycle |__| Bicycle |__| Computer |__| Refrigerator |__| Microwave |__| Television |__| Radio |__|

HIV Knowledge Questionnaire 18: For each statement, please tick "True", "False" or "I don't know" (DK). If you do not know, please do not guess; instead tick "DK"

- | | | | | | | |
|--|------|----|-------|----|----|----|
| 1. Coughing and sneezing DO NOT spread HIV? | True | __ | False | __ | DK | __ |
| 2. A person can get HIV by sharing a glass of water with someone who has HIV? | True | __ | False | __ | DK | __ |
| 3. Pulling out the penis before a man climaxes/cams keeps a woman from getting HIV during sex | True | __ | False | __ | DK | __ |
| 4. A woman can get HIV if she has anal sex with a man | True | __ | False | __ | DK | __ |
| 5. Showering, or washing one's genitals/private parts, after sex keeps a person from getting HIV | True | __ | False | __ | DK | __ |
| 6. All pregnant women infected with HIV will have babies born with AIDS | True | __ | False | __ | DK | __ |
| 7. People who have been infected with HIV quickly show serious signs of being infected | True | __ | False | __ | DK | __ |
| 8. There is a vaccine that can stop adults from getting HIV | True | __ | False | __ | DK | __ |
| 9. People are likely to get HIV by deep kissing, putting their tongue in their partner's mouth, if their partner has HIV | True | __ | False | __ | DK | __ |

- | | | | | | | |
|---|------|--------------------------|-------|--------------------------|----|--------------------------|
| 10. A woman can get HIV if she has sex during her period | True | <input type="checkbox"/> | False | <input type="checkbox"/> | DK | <input type="checkbox"/> |
| 11. There is a female condom that can help decrease a woman's chance of getting HIV | True | <input type="checkbox"/> | False | <input type="checkbox"/> | DK | <input type="checkbox"/> |
| 12. A natural skin condom works better against HIV than does a latex condom | True | <input type="checkbox"/> | False | <input type="checkbox"/> | DK | <input type="checkbox"/> |
| 13. A person will NOT get HIV if she or he is taking antibiotics | True | <input type="checkbox"/> | False | <input type="checkbox"/> | DK | <input type="checkbox"/> |
| 14. Having sex with more than one partner can increase a person's chance of being infected with HIV | True | <input type="checkbox"/> | False | <input type="checkbox"/> | DK | <input type="checkbox"/> |
| 15. Taking a test for HIV one week after having sex will tell a person if she or he has HIV | True | <input type="checkbox"/> | False | <input type="checkbox"/> | DK | <input type="checkbox"/> |
| 16. A person can get HIV by sitting in a hot tub or a swimming pool with a person who has HIV | True | <input type="checkbox"/> | False | <input type="checkbox"/> | DK | <input type="checkbox"/> |
| 17. A person can get HIV from oral sex | True | <input type="checkbox"/> | False | <input type="checkbox"/> | DK | <input type="checkbox"/> |
| 18. Using Vaseline or baby oil with condoms lowers the chance of getting HIV | True | <input type="checkbox"/> | False | <input type="checkbox"/> | DK | <input type="checkbox"/> |

SEXUAL AND HIV PREVENTIVE BEHAVIOR: Let me assure you again that your answers are completely confidential and will not be told to anyone. If we should come to any question that you do not want to answer, just let me know and we will go on to the next question.

1. When was the last time you had sexual intercourse (record days/weeks/months)? | Prefer not to answer Never
2. The last time you had sexual intercourse, was a condom used? Yes No Prefer not to answer
3. What was your relationship with the person you last had sex with? Live-in partner Boyfriend/girlfriend, not live in Casual acquaintance Prostitute Other | Prefer not to answer
4. The last time you had intercourse, did you or your partner drink alcohol? Yes No Prefer not to answer
5. If yes to (4), were you or your partner drunk? Respondent Partner Both Prefer not to answer
6. How many different people have you had sex with since the last interview? | Prefer not to answer
7. Recall the time since our last interview; did you use a condom every time you had sex? always only sometimes rarely never Prefer not to answer
8. Did you pay (or get paid) in exchange for sex? Yes No Prefer not to answer
9. Do you know your HIV status? Yes No Prefer not to answer
10. Have you tested for HIV since our last interview? Yes No Prefer not to answer
11. Are you circumcised? Yes No Prefer not to answer
12. If no, would you consider getting circumcised? Yes No Prefer not to answer
13. Do you know where you can get condoms? Yes No Prefer not to answer
14. If YES, probe for place(s)
15. Would you yourself go to acquire a condom if you needed one? Yes No Prefer not to answer